

A purification postulate for quantum mechanics with indefinite causal order

Mateus Araújo^{1 2}, Adrien Feix^{1 2}, Miguel Navascués^{1 2}, and Časlav Brukner^{1 2}

¹Faculty of Physics, University of Vienna, Boltzmanngasse 5 1090 Vienna, Austria

²Institute for Quantum Optics and Quantum Information (IQOQI), Boltzmanngasse 3 1090 Vienna, Austria

29th November 2016

To study which are the most general causal structures which are compatible with local quantum mechanics, Oreshkov *et al.* [1] introduced the notion of a process: a resource shared between some parties that allows for quantum communication between them without a predetermined causal order. These processes can be used to perform several tasks that are impossible in standard quantum mechanics: they allow for the violation of causal inequalities, and provide an advantage for computational and communication complexity. Nonetheless, no process that can be used to violate a causal inequality is known to be physically implementable. There is therefore considerable interest in determining which processes are physical and which are just mathematical artefacts of the framework. Here we make the first step in this direction, by proposing a purification postulate: processes are physical only if they are purifiable. We derive necessary conditions for a process to be purifiable, and show that several known processes do not satisfy them.

1 Introduction

It is widely believed that any future theory that unifies quantum mechanics and gravity will feature quantum uncertainty in the metric tensor [2], which should produce indefinite causal structures. Our understanding of the notion of indefinite causal structures is, however, still lacking. To investigate that, one approach is to consider simple, concrete models that are compatible with the laws of quantum information processing. One such model – the process matrix formalism – was introduced by Oreshkov *et al.* as the most general causal structure that can connect local laboratories where standard quantum mechanics is valid without creating paradoxical causal loops [1].

These processes have been shown to enable the realization of tasks that are otherwise impossible: they allow for the violation of causal inequalities [1, 3–8], can be detected by causal witnesses [9], provide an advantage in quantum computation [10–12], and enable a reduction in communication complexity [13, 14]. But even though one such process – the quantum switch – has been implemented experimentally [15, 16], in general it is not known if all process are physical or some of them are just mathematical artefacts of the formalism. They were, after all, defined only from the requirement of not generating logical contradictions, and processes realisable in nature are likely to fulfil additional physical constraints.

One can, therefore, look for requirements beyond mere logical consistency in order to shed light on the physicality of these processes. As seen from the search of physical principles to determine the set of quantum correlations, such meta-theoretical principles can provide nontrivial constraints on the possible theories [17–21]. The principle we choose to investigate here is reversibility of the transformations between quantum states: it is, after all, valid in all fundamental physical theories, and was a central ingredient in all of the reconstructions of quantum mechanics [22–28].

To define what reversibility means for processes we first needed to generalise their definition: we extend them with a global past and a global future, so that they can be seen as inducing a transformation from quantum states in the past to quantum states in the future, after passing through the causally indefinite region of the local laboratories. We can then define pure processes as those that preserve the reversibility of the underlying operations, i.e., those that induce a unitary transformation from the past to the future whenever unitary transformations are also applied in the local laboratories.

With these definitions in hand, we can propose the purification postulate: *processes are physical only if they are purifiable*, i.e., if they can be expressed as a part of a pure process in a larger space. It turns out to be rather difficult to determine if a given process is purifiable. We derive, then, a necessary but not sufficient condition

for purifiability, and show that several known processes fail to satisfy this condition, among which the process from Ref. [1] that was the first one shown to violate a causal inequality. In fact, no bipartite process that was able to violate a causal inequality was found to be purifiable. There exists, however, a tripartite process which is purifiable and able to violate causal inequalities. This implies that the purification postulate is necessary but not sufficient for physicality of a process, if one takes the view that violations of causal inequalities is impossible in Nature [29].

The paper is organized as follows: in section 2 we generalise the definition of a process and introduce the notion of pure processes. In section 3 we propose the purification postulate. In section 4 we derive necessary and sufficient conditions for a process to be purifiable, and in section 5 we derive a necessary but not sufficient condition for purifiability that can be easily tested. In section 6 we apply our condition to several known processes, and in section 7 we present a tripartite pure process that violates causal inequalities.

2 Pure processes

In Ref. [1], the process matrix was introduced as a (multi)linear function that takes completely positive (CP) maps to probabilities. To be able to talk about purification, we need to extend this definition to take CP maps not to probabilities, but to other CP maps. This view of a process as a higher-order transformation is much in the spirit of quantum combs [30], and can be recovered from the multipartite definition of process presented for example in Ref. [9]. We shall, nevertheless, explicitly define processes as higher-order transformations for clarity and to make this paper self-contained.

To this end, let \mathcal{A} be a completely positive trace preserving (CPTP) map that takes the Hilbert spaces A_I, A'_I to A_O, A'_O , and \mathcal{B} a CPTP map that takes B_I, B'_I to B_O, B'_O . A process is then defined as the most general linear transformation that acts trivially on A'_I, B'_I, A'_O, B'_O and takes \mathcal{A} and \mathcal{B} to a CPTP map $\mathcal{G}_{\mathcal{A},\mathcal{B}}$ from A'_I, B'_I, P to A'_O, B'_O, F , as represented in Fig. 1. Note that the Hilbert space F cannot signal to any other Hilbert space, and can therefore be interpreted as a global future. Conversely, none of the Hilbert spaces can signal to P , which can then be interpreted as a global past.




Figure 1: A process W is a bilinear function from the CPTP maps \mathcal{A}, \mathcal{B} to a CPTP map $\mathcal{G}_{\mathcal{A},\mathcal{B}}$. This map takes states from the global past P and auxiliary spaces A'_I, B'_I to the global future F and auxiliary spaces A'_O, B'_O .

To find a matrix representation for a process, we make use of the Choi-Jamiołkowski (CJ) isomorphism, which we recap in appendix A. Let then $A = \mathfrak{C}(\mathcal{A})$, $B = \mathfrak{C}(\mathcal{B})$, and $G_{\mathcal{A},\mathcal{B}} = \mathfrak{C}(\mathcal{G}_{\mathcal{A},\mathcal{B}})$ be the CJ representations of the CPTP maps \mathcal{A} , \mathcal{B} , and $\mathcal{G}_{\mathcal{A},\mathcal{B}}$. By linearity we can represent the mapping from A, B to $G_{\mathcal{A},\mathcal{B}}$ as

$$G_{\mathcal{A},\mathcal{B}} = \text{tr}_{A_I A_O B_I B_O} [W^{T_{A_I A_O B_I B_O}} (A \otimes B)], \quad (1)$$

where $W \in P \otimes F \otimes A_I \otimes A_O \otimes B_I \otimes B_O$ is the process matrix we are defining, and identity matrices on the subsystems P , F , A'_I , A'_O , B'_I , and B'_O are left implicit. This formula can be rewritten in a more convenient way using the link product, which was designed to conveniently express the CJ representation of a composition of quantum operations [30], and we recap in appendix B:

$$G_{\mathcal{A},\mathcal{B}} = W * (A \otimes B) \quad (2)$$

We need $G_{\mathcal{A},\mathcal{B}}$ to be a valid CPTP map for all valid CPTP maps A and B . This imposes the following restrictions on W , which we derive in appendix C:

$$W \geq 0, \quad (3)$$

$$\text{tr } W = d_{A_O} d_{B_O} d_P, \quad (4)$$

$$W = L_V(W), \quad (5)$$

where L_V is a projector on the linear subspace of valid process matrices defined in equation (52).

As an explicit example of a process defined as a higher-order transformation consider a situation where a state in $P = P_1 \otimes P_2$ is sent to Alice and Bob, and the resulting state from Alice and Bob is sent to $F = F_1 \otimes F_2$. The process matrix is given by $|w_{\text{state}}\rangle\langle w_{\text{state}}|$, where

$$|w_{\text{state}}\rangle = |\mathbb{1}\rangle^{P_1 A_I} |\mathbb{1}\rangle^{P_2 B_I} |\mathbb{1}\rangle^{A_O F_1} |\mathbb{1}\rangle^{B_O F_2}, \quad (6)$$

and $|\mathbb{1}\rangle$ is the “pure” CJ representation of the identity map between the appropriate Hilbert spaces, as defined in Appendix A.

A less trivial example is the process where a state in P goes first to Alice, then to Bob, and finally to F . Using again the vector representation the process is

$$|w_{\text{channel}}\rangle = |\mathbb{1}\rangle^{P A_I} |\mathbb{1}\rangle^{A_O B_I} |\mathbb{1}\rangle^{B_O F}. \quad (7)$$

Finally, an example of a process that encodes an indefinite causal order is the quantum switch [9, 10], which in this representation is:

$$|w_{\text{switch}}\rangle = |0\rangle^{P_1} |\mathbb{1}\rangle^{P_2 A_I} |\mathbb{1}\rangle^{A_O B_I} |\mathbb{1}\rangle^{B_O F_2} |0\rangle^{F_1} + |1\rangle^{P_1} |\mathbb{1}\rangle^{P_2 B_I} |\mathbb{1}\rangle^{B_O A_I} |\mathbb{1}\rangle^{A_O F_2} |1\rangle^{F_1} \quad (8)$$

To connect this version of the quantum switch with the one in Refs. [9, 10], one should prepare the control qubit and send it to P_1 , and prepare the target qubit and send it to P_2 . After the process is done the control qubit will be found in F_1 (unchanged), and the target qubit in F_2 .

This definition of processes allows a natural definition of what it means for a process to be “pure”. Just as we can define unitaries as the most general linear transformations that map pure states to pure states of the same dimension¹, we can define pure processes to be most general linear transformations that map unitaries to unitaries. More precisely:

Definition 1. A process W is pure if for all unitaries \mathcal{A}, \mathcal{B} the resulting transformation $\mathcal{G}_{\mathcal{A}, \mathcal{B}}$ is also a unitary.

It turns out that pure processes are unitary transformations from A_O, B_O, P to A_I, B_I, F and can be conveniently represented as vectors, as shown in the following theorem:

Theorem 2. A process W is pure if and only if $W = |U_w\rangle\langle U_w|$ for some unitary U_w .

Proof. If W is pure, then $\mathcal{G}_{\mathcal{A}, \mathcal{B}}$ must be a unitary in particular when \mathcal{A} and \mathcal{B} are SWAPs that map A'_I, A_I to A_O, A'_O and B'_I, B_I to B_O, B'_O . Then

$$G_{\mathcal{A}, \mathcal{B}} = W * \left(|\mathbb{1}\rangle\langle\mathbb{1}|^{A'_I A_O} |\mathbb{1}\rangle\langle\mathbb{1}|^{A_I A'_O} \otimes |\mathbb{1}\rangle\langle\mathbb{1}|^{B'_I B_O} |\mathbb{1}\rangle\langle\mathbb{1}|^{B_I B'_O} \right) = W, \quad (9)$$

so the resulting transformation $\mathcal{G}_{\mathcal{A}, \mathcal{B}} = \mathfrak{C}^{-1}(G_{\mathcal{A}, \mathcal{B}}) = \mathfrak{C}^{-1}(W) =: \mathcal{W}$ is just the process itself, with the relabelling $A_I \mapsto A'_O, A_O \mapsto A'_I, B_I \mapsto B'_O$, and $B_O \mapsto B'_I$, so \mathcal{W} must be a unitary transformation from A_O, B_O, P to A_I, B_I, F . Writing $\mathcal{W}(\rho) = U_w \rho U_w^\dagger$, we have that its CJ representation is $W = \mathfrak{C}(\mathcal{W}) = |U_w\rangle\langle U_w|$.

Conversely, if $W = |U_w\rangle\langle U_w|$, $A = |U_a\rangle\langle U_a|$, and $B = |U_b\rangle\langle U_b|$, then

$$G_{\mathcal{A}, \mathcal{B}} = |U_w\rangle\langle U_w| * (|U_a\rangle\langle U_a| \otimes |U_b\rangle\langle U_b|) = |U_g\rangle\langle U_g|, \quad (10)$$

where

$$U_g^{A'_I B'_I P} = \text{tr}_{A_I B_I} \left[\left(U_w^{P A_I B_I} \otimes \mathbb{1}^{A'_I B'_I} \right) \left(\mathbb{1}^P \otimes U_a^{A_I A'_I} \otimes U_b^{B_I B'_I} \right) \right]. \quad (11)$$

Since by assumption $\mathcal{G}_{\mathcal{A}, \mathcal{B}}$ is trace preserving, we have that $\text{tr}_{A'_O B'_O F} G_{\mathcal{A}, \mathcal{B}} = \mathbb{1}^{A'_I B'_I P}$. Substituting $G_{\mathcal{A}, \mathcal{B}} = |U_g\rangle\langle U_g|$ we get

$$\text{tr}_{A'_O B'_O F} |U_g\rangle\langle U_g| = U_g^\dagger U_g = \mathbb{1}^{A'_I B'_I P}. \quad (12)$$

Since, furthermore, U_g is a square finite-dimensional matrix, it has a right inverse which is equal to its left inverse, so U_g is a unitary, and so is $\mathcal{G}_{\mathcal{A}, \mathcal{B}}$. \square

¹More precisely, a linear map \mathcal{E} is a unitary iff for all pure states $|\psi\rangle$ the transformed state $\mathcal{I} \otimes \mathcal{E}(|\psi\rangle\langle\psi|)$ is a pure state of the same dimension.

3 A purification postulate

With the definition of a pure process in hand, we can now define what it means to purify a process: we say that a process W with trivial P and F is purifiable if one can recover it from a pure process S by inputting the state² $|0\rangle$ in P and tracing out F , i.e., if

$$W = S * (|0\rangle\langle 0|^P \otimes \mathbb{1}^F), \quad (13)$$

as shown in Fig. 2.




Figure 2: A process W with trivial P and F is purifiable if it can be recovered from a pure process S by inputting the state $|0\rangle$ in P and tracing out F .

We propose then the purification postulate: *a process is physical only if it is purifiable*. This postulate is motivated by the fact that a non-purifiable process would fundamentally map unitaries onto isometries or non-unitary CPTP maps, destroying the reversibility of the theory. Reversibility, on its turn, is a cherished principle: all fundamental physical theories are time reversible³ and the hint that the formation and evaporation of a black hole might be an irreversible evolution is one of the major problems in modern physics [34, 35]. Moreover, in all reconstructions of quantum mechanics [22–28] reversibility was used as a central ingredient. This supports the idea that it is indeed a fundamental part of quantum mechanics, and it should not be done away with lightly. Furthermore, although there is speculation that in a full theory of quantum gravity new degrees of freedom are created by the expansion of the Universe [36–38], in concrete models these degrees of freedom are born in the vacuum state [39], making the time evolution actually be reversible. In any case, current inflationary models using quantum field theory on a curved expanding spacetime do not feature creation of new degrees of freedom [40].

As a sanity check, note that all processes known to be physical – all processes where the order between the parties is fixed, e.g. $|w_{\text{channel}}\rangle$ (equation (7)), controlled by a quantum system, e.g. $|w_{\text{switch}}\rangle$ (equation (8)), or incoherently mixed – are purifiable.

4 Necessary and sufficient conditions for purification

We shall now derive necessary and sufficient conditions for a process W with trivial P and F to be purifiable, i.e., to be recoverable via equation (13) from a process S that is pure and valid. From Theorem 2, we know that S is pure iff $S = |U_s\rangle\rangle\langle\langle U_s|$ for some unitary U_s . Defining

$$|w_\psi\rangle := \langle\psi^*|^P |U_s\rangle\rangle \quad (14)$$

and noting that

$$|U_s\rangle\rangle\langle\langle U_s| * (|\psi\rangle\langle\psi|^P \otimes \mathbb{1}^F) = |w_\psi\rangle\langle w_\psi| * \mathbb{1}^F = \text{tr}_F |w_\psi\rangle\langle w_\psi|, \quad (15)$$

we can rewrite equation (13) as

$$W = \text{tr}_F |w_0\rangle\langle w_0|, \quad (16)$$

where $|w_0\rangle = \langle 0|^P |U_s\rangle\rangle$. We can also state the condition that S is valid purely in terms of the vectors $|w_\psi\rangle$. To do that, remember that S is valid iff $S * (A \otimes B)$ is a CPTP map for all CPTP maps A and B . In its turn,

²We can fix the state to be $|0\rangle$ instead of allowing an arbitrary state without loss of generality.

³Note that to treat the collapse of the wavefunction during a measurement as an irreversible process one needs to use objective collapse models [31–33] instead of standard quantum mechanics. We are taking the view that collapse is not a physical process.

$S * (A \otimes B)$ is a CPTP map iff for all input states $|\psi\rangle$ its output $[S * (A \otimes B)] * |\psi\rangle\langle\psi|^P$ is a valid quantum state. Note, however, that

$$[S * (A \otimes B)] * |\psi\rangle\langle\psi|^P = (S * |\psi\rangle\langle\psi|^P) * (A \otimes B),$$

and the condition that the right hand side is a valid quantum state for all $|\psi\rangle$, A , and B is precisely the condition that the process with trivial P given by $(S * |\psi\rangle\langle\psi|^P)$ produces a valid CPTP (with trivial P) when linked with the CPTP maps A and B , which means that the process $(S * |\psi\rangle\langle\psi|^P) = |w_\psi\rangle\langle w_\psi|$ is a valid process. Writing out the validity conditions explicitly, we have

$$\forall |\psi\rangle \quad |w_\psi\rangle\langle w_\psi| = L_V(|w_\psi\rangle\langle w_\psi|) \quad \text{and} \quad \text{tr}|w_\psi\rangle\langle w_\psi| = d_{A_O} d_{B_O}, \quad (17)$$

where L_V is the projector onto the valid subspace defined in equation (52) particularized for $d_P = 1$.

With this, we reduce the problem of purifying a process W to finding a set of vectors $|w_\psi\rangle$ such that equations (16) and (17) are satisfied. We can simplify it further by noting that equation (16) is just the purification of a positive matrix, which admits a simple solution. Let then the eigendecomposition of W be

$$W = \sum_{i=0}^{r-1} \lambda_i |\lambda_i\rangle\langle\lambda_i|, \quad (18)$$

where r is the rank of W , and $\lambda_i, |\lambda_i\rangle$ its nonzero eigenvalues and corresponding eigenvectors. Then a valid purification for it is

$$|w_0\rangle = \sum_{i=0}^{r-1} \sqrt{\lambda_i} |\lambda_i\rangle |i\rangle, \quad (19)$$

which is unique modulo some isometry V on the purifying system. But this isometry has no effect on the validity of $|w_\psi\rangle\langle w_\psi|$, as it only affects the state output by the process, and an isometry maps valid quantum states to valid quantum states. This implies that we can choose without loss of generality the isometry to be the identity, and the dimension of the purifying system $d_F = d_P$ to be the rank of W . This allows us to use a r^2 -dimensional basis for P to simplify condition (17). Choosing $\{|i\rangle\langle j|\}_{i,j=0}^{r-1}$ as this basis, and rewriting the equations explicitly in terms of the vectors $|w_i\rangle = \langle i|^P |U_s\rangle$, the condition translates to

$$\forall i, j \quad L_V^\perp(|w_i\rangle\langle w_j|) = 0 \quad \text{and} \quad \langle w_i | w_j \rangle = d_{A_O} d_{B_O} \delta_{ij},$$

where for brevity we are using the projector $L_V^\perp := \mathbb{1} - L_V$.

We summarize the results of this section in the following theorem:

Theorem 3. *A process matrix W of rank r and eigendecomposition*

$$W = \sum_{i=0}^{r-1} \lambda_i |\lambda_i\rangle\langle\lambda_i| \quad (20)$$

is purifiable if and only if there exists a set of vectors $\{|w_i\rangle\}_{i=0}^{r-1}$ such that

$$|w_0\rangle = \sum_{i=0}^{r-1} \sqrt{\lambda_i} |\lambda_i\rangle |i\rangle \quad (21)$$

and

$$\forall i, j \quad L_V^\perp(|w_i\rangle\langle w_j|) = 0 \quad \text{and} \quad \langle w_i | w_j \rangle = d_{A_O} d_{B_O} \delta_{ij} \quad (22)$$

If they exist, a process $S = |U_s\rangle\langle\langle U_s|$ that purifies W is given by

$$\langle i|^P |U_s\rangle := |w_i\rangle \quad (23)$$

5 Necessary condition for purification

It is not easy to find a set of vectors $\{|v_i\rangle\}$ that satisfies the conditions of Theorem 3, as the constraint $L_V^\perp(|v_i\rangle\langle v_j|) = 0$ is quadratic on them. We shall not solve this problem in full, but rather prove an upper bound on the number of vectors that can satisfy those conditions for a given process. If this upper bound is smaller than the rank of the process, this is a proof that this process is not purifiable.

We shall start by focussing in the simplest of the conditions, namely that $L_V^\perp(|v_i\rangle\langle w_0|) = 0$. Since this condition is linear on $|v_i\rangle$ it is straightforward to characterize the vector space V_W formed by the vectors that satisfy it. Restricting our attention to this (hopefully small) subspace makes it easier to consider the other, non-linear, conditions.

To construct an orthonormal basis for V_W , first we transform the projector L_V^\perp to act on the “double-kets” of process matrices, i.e., we define the matrix $\Pi_{L_V^\perp}$ such that⁴

$$\forall i, j \quad \Pi_{L_V^\perp} |i\rangle\langle j| \rangle\rangle = |L_V^\perp(|i\rangle\langle j|) \rangle\rangle$$

We have then

$$L_V^\perp(|v\rangle\langle w_0|) = 0 \iff \Pi_{L_V^\perp} |w_0^*\rangle|v\rangle = 0,$$

so if we define

$$O_W := (\langle w_0^*| \otimes \mathbb{1}) \Pi_{L_V^\perp} (|w_0^*\rangle \otimes \mathbb{1})$$

we have that

$$O_W |v\rangle = 0 \iff \Pi_{L_V^\perp} |w_0^*\rangle|v\rangle = 0,$$

so the null eigenvectors of O_W form an orthonormal basis for V_W . With it in hand we can now restrict the projector $\Pi_{L_V^\perp}$ to the subspace⁵ $V_W^* \otimes V_W$. Let $\{|r_i\rangle\}$ be an orthonormal basis for V_W , and $R = \sum_i |i\rangle\langle r_i|$. The restriction $\Pi_{L_V^\perp|V_W}$ is then given by

$$\Pi_{L_V^\perp|V_W} = (R^* \otimes R) \Pi_{L_V^\perp} (R^{\dagger} \otimes R^{\dagger}).$$

We can now consider the condition that $L_V^\perp(|v_i\rangle\langle v_j|) = 0$. Let $\{|m_k\rangle\}_{k=0}^{d_m-1}$ be the set of eigenvectors of $\Pi_{L_V^\perp|V_W}$ with nonzero eigenvalue⁶. Then

$$\Pi_{L_V^\perp|V_W} |v_j^*\rangle|v_i\rangle = 0 \iff \forall k \quad \langle v_j^*| \langle v_i| |m_k\rangle = 0.$$

We want to rewrite the inner product $\langle v_j^*| \langle v_i| |m_k\rangle$ in a more convenient way. For that, note that $\langle \psi| A | \psi\rangle = \langle \psi^*| \langle \psi| A^T \rangle\rangle$, so $\langle v_j^*| \langle v_i| |m_k\rangle = \langle v_i| M_k | v_j\rangle$ for matrices M_k such that $|M_k^T\rangle\rangle = |m_k\rangle$. These matrices are in general not Hermitian, so for convenience we define $C_k = M_k + M_k^\dagger$ and $C_{k+d_m} = i(M_k - M_k^\dagger)$. Then

$$\Pi_{L_V^\perp|V_W} |v_j^*\rangle|v_i\rangle = 0 \iff \forall k \quad \langle v_i| C_k | v_j\rangle = 0.$$

Let now d_{C_k} be the dimension of the largest subspace such that all $|v\rangle$ in this subspace respect $\langle v| C_k | v\rangle = 0$. We can easily calculate d_{C_k} using the the null-square lemma proved in Appendix D: let n_{k+} , n_{k-} , and n_{k0} be the number of positive, negative, and null eigenvalues of C_k . Then

$$d_{C_k} = n_{k0} + \min\{n_{k+}, n_{k-}\},$$

and a simple upper bound on the dimension of the largest subspace of vectors that respect the conditions of Theorem 3 is

$$d_{\max}(W) := \min_i d_{C_i} \tag{24}$$

Therefore if $d_{\max}(W) < \text{rank}(W)$ the process W is not purifiable.

6 Examples

We shall now apply the necessary condition derived in the previous section to several process matrices from the literature. In this section we shall omit superscripts that identify subsystems and tensor products to avoid clutter. The expression (e.g.) $Z\mathbb{1}XZ$ should be understood as $Z^{A_I} \otimes \mathbb{1}^{A_O} \otimes X^{B_I} \otimes Z^{B_O}$.

Let

$$W_{\text{OCB}} = \frac{1}{4} \left[\mathbb{1}\mathbb{1}\mathbb{1}\mathbb{1} + \frac{\mathbb{1}ZZ\mathbb{1} + Z\mathbb{1}XZ}{\sqrt{2}} \right] \tag{25}$$

⁴ $\Pi_{L_V^\perp}$ can be explicitly written as $\Pi_{L_V^\perp} = \mathfrak{C}(L_V^\perp)^R$, where R is the reshuffling operation [41].

⁵One can restrict instead to $V_W \setminus |w_0\rangle$ for a slight computational advantage.

⁶It is possible that this set is empty. In this case all vectors in V_W respect the conditions of Theorem 3, and the process is purifiable iff $\dim(V_W) \geq \text{rank}(W)$.

be the process introduced in Ref. [1]. It was proven, under a restriction on the allowed operations, to produce the maximal violation of the original causal inequality for any dimension [42]. Since $\text{rank}(W_{\text{OCB}}) = 8$ and $d_{\text{max}}(W_{\text{OCB}}) = 5$ it is not purifiable.

Let

$$W_{\text{max}} = \frac{1}{4} \left[\mathbb{1}\mathbb{1}\mathbb{1}\mathbb{1} + a_0 Z\mathbb{1}Z\mathbb{1} - a_1 (Z\mathbb{1}\mathbb{1}\mathbb{1} + \mathbb{1}\mathbb{1}Z\mathbb{1}) - a_2 (Z\mathbb{1}\mathbb{1}Z + \mathbb{1}ZZ\mathbb{1}) + a_3 (Z\mathbb{1}ZZ + ZZZ\mathbb{1}) + a_4 (Z\mathbb{1}XX - Z\mathbb{1}YY + XXZ\mathbb{1} - YYZ\mathbb{1}) \right], \quad (26)$$

where $a_0 \approx 0.2744$, $a_1 \approx 0.2178$, $a_2 \approx 0.3628$, $a_3 \approx 0.3114$, and $a_4 \approx 0.2097$. This process was introduced in Appendix C of Ref. [5] and conjectured to produce the maximal violation of GYNI and LGYNI inequalities for its dimension. Since $\text{rank}(W_{\text{max}}) = 11$ and $d_{\text{min}}(W_{\text{max}}) = 10$ it is not purifiable.

Let

$$W_{\text{opt}} = \frac{1}{4} \left[\mathbb{1}\mathbb{1}\mathbb{1}\mathbb{1} + \frac{1}{\sqrt{3}} Z\mathbb{1}XZ + \frac{\sqrt{3}-1}{3} (\mathbb{1}XX\mathbb{1} + \mathbb{1}YY\mathbb{1} + \mathbb{1}ZZ\mathbb{1}) \right] \quad (27)$$

be the process introduced in Ref. [29]. By itself it cannot violate any causal inequalities, but it becomes able to do so when it is extended with an entangled ancilla. However, when admixed with a small amount of noise, the known violations disappear, so this noisy version was conjectured to be unable to violate any causal inequalities. That is, W_{opt} was conjectured to be “extensibly causal”. Since extensibly causal processes seem physically reasonable, but we have no physical interpretation for W_{opt} or its noisy version, it would very interesting to test its purifiability to gather evidence about its physicality. Unfortunately, since $\text{rank}(W_{\text{opt}}) = 12$ and $d_{\text{max}}(W_{\text{opt}}) = 17$ our test is inconclusive. This is also the case for the noisy version, which has rank 16 and d_{max} equal to 41.

7 Counter-example

Since the processes examined in the previous section either are not purifiable or have unknown status, it might be tempting to conjecture that all processes that can violate causal inequalities are not purifiable. While this might be true for the bipartite case, there exists a counter-example for the general case: a tripartite process which can violate causal inequalities and is purifiable.

This process, which was first published in Ref. [6], can be concisely described as a function from A_O, B_O, C_O to A_I, B_I, C_I that incoherently maps the basis states $|a, b, c\rangle$ to the basis states $|\neg b \wedge c, \neg c \wedge a, \neg a \wedge b\rangle$, where \neg and \wedge represent logical negation and logical conjunction. It can be represented in a more cumbersome way as the process matrix

$$W_{\text{det}} = \sum_{abc} |a, b, c\rangle\langle a, b, c| \otimes |\neg b \wedge c, \neg c \wedge a, \neg a \wedge b\rangle\langle \neg b \wedge c, \neg c \wedge a, \neg a \wedge b| \quad (28)$$

where unlike in the other processes we wrote the subsystems in the order $A_O, B_O, C_O, A_I, B_I, C_I$ to match the function described above. To purify this process [43] we use the standard trick to turn a irreversible function into a reversible one, that is, changing the function from $|x\rangle \mapsto |f(x)\rangle$ to $|x\rangle|y\rangle \mapsto |x\rangle|y \oplus f(x)\rangle$. In terms of our function this means mapping $|a, b, c\rangle|i, j, k\rangle$ to $|a, b, c\rangle|i \oplus \neg b \wedge c, j \oplus \neg c \wedge a, k \oplus \neg a \wedge b\rangle$. This reversible function is then the purification of W_{det} , and can be written as the process vector

$$|w_{\text{det}}\rangle = \sum_{\substack{abc \\ ijk}} |a, b, c\rangle|i, j, k\rangle \otimes |a, b, c\rangle|i \oplus \neg b \wedge c, j \oplus \neg c \wedge a, k \oplus \neg a \wedge b\rangle \quad (29)$$

where the subsystems are written in the order $A_O, B_O, C_O, P, F, A_I, B_I, C_I$. Note that unlike the other subsystems P and F consist of three qubits each.

8 Conclusion

We have proposed a purification postulate, motivated by considerations that are independent of this formalism, that allows us for the first time to declare a large class of processes to be unphysical. This can be made part of a axiomatic, top-down approach to determining the set of physical processes. Ideally, one should marry it with a bottom-up approach, that shows processes to be physical by actually constructing laboratory implementations for them. As of yet, however, there are still processes which are neither excluded by the purification principle nor known to be implementable, showing that more work must be done in order to consummate this marriage.

In this gray zone we find, for example, $|w_{\text{det}}\rangle$ (equation (29)), showing that the set of purifiable processes is rather rich, and that the purification principle does not rule out the violation of causal inequalities in Nature. It might still be the case, however, that processes that cannot violate causal inequalities, i.e., extensively causal processes, are always purifiable. In order to decide this it would be interesting to know whether the noisy version of W_{opt} (equation (27)) is purifiable.

9 Acknowledgments

We thank Āmin Baumeler, Fabio Costa, Paolo Perinotti, and Stefan Wolf for useful discussions. We acknowledge support from the European Commission project RAQUEL (No. 323970); the Austrian Science Fund (FWF) through the Special Research Programme FoQuS, the Doctoral Programme CoQuS and Individual Project (No. 24621), and the John Templeton Foundation.

References

- [1] O. Oreshkov, F. Costa and Ā. Brukner. ‘Quantum correlations with no causal order’. *Nat. Commun.* **3** 1092 (2012). [arXiv:1105.4464 \[quant-ph\]](#).
- [2] A. Ashtekar. ‘Large Quantum Gravity Effects: Unforeseen Limitations of the Classical Theory’. *Phys. Rev. Lett.* **77** 4864–4867 (1996). [arXiv:gr-qc/9610008](#).
- [3] Ā. Baumeler and S. Wolf. ‘Perfect signaling among three parties violating predefined causal order’. *Information Theory (ISIT), 2014 IEEE International Symposium on* 526–530 (2014). [arXiv:1312.5916 \[quant-ph\]](#).
- [4] Ā. Baumeler, A. Feix and S. Wolf. ‘Maximal incompatibility of locally classical behavior and global causal order in multi-party scenarios’. *Phys. Rev. A* **90** 042106 (2014). [arXiv:1403.7333 \[quant-ph\]](#).
- [5] C. Branciard, M. Araújo, A. Feix, F. Costa and Ā. Brukner. ‘The simplest causal inequalities and their violation’. *New J. Phys.* **18** 013008 (2015). [arXiv:1508.01704 \[quant-ph\]](#).
- [6] Ā. Baumeler and S. Wolf. ‘The space of logically consistent classical processes without causal order’. *New J. Phys.* **18** 013036 (2016). [arXiv:1507.01714 \[quant-ph\]](#).
- [7] O. Oreshkov and C. Giarmatzi. ‘Causal and causally separable processes’. *New J. Phys.* **18** 093020 (2015). [arXiv:1506.05449 \[quant-ph\]](#).
- [8] A. A. Abbott, C. Giarmatzi, F. Costa and C. Branciard. ‘Multipartite Causal Correlations: Polytopes and Inequalities’ (2016). [arXiv:1608.01528 \[quant-ph\]](#).
- [9] M. Araújo, C. Branciard, F. Costa, A. Feix, C. Giarmatzi and Āslav Brukner. ‘Witnessing causal non-separability’. *New J. Phys.* **17** 102001 (2015). [arXiv:1506.03776 \[quant-ph\]](#).
- [10] G. Chiribella, G. M. D’Ariano, P. Perinotti and B. Valiron. ‘Quantum computations without definite causal structure’. *Phys. Rev. A* **88** 022318 (2013). [arXiv:0912.0195 \[quant-ph\]](#).
- [11] G. Chiribella. ‘Perfect discrimination of no-signalling channels via quantum superposition of causal structures’. *Phys. Rev. A* **86** 040301 (2012). [arXiv:1109.5154 \[quant-ph\]](#).
- [12] M. Araújo, F. Costa and Ā. Brukner. ‘Computational Advantage from Quantum-Controlled Ordering of Gates’. *Phys. Rev. Lett.* **113** 250402 (2014). [arXiv:1401.8127 \[quant-ph\]](#).
- [13] A. Feix, M. Araújo and Ā. Brukner. ‘Quantum superposition of the order of parties as a communication resource’. *Phys. Rev. A* **92** 052326 (2015). [arXiv:1508.07840 \[quant-ph\]](#).
- [14] P. Allard Guérin, A. Feix, M. Araújo and Ā. Brukner. ‘Exponential communication complexity advantage from quantum superposition of the direction of communication’. *Phys. Rev. Lett.* **117** 100502 (2016). [arXiv:1605.07372 \[quant-ph\]](#).
- [15] L. M. Procopio, A. Moqanaki, M. Araújo, F. Costa, I. A. Calafell, E. G. Dowd, D. R. Hamel, L. A. Rozema, Ā. Brukner and P. Walther. ‘Experimental superposition of orders of quantum gates’. *Nat. Commun.* **6** 7913 (2015). [arXiv:1412.4006 \[quant-ph\]](#).
- [16] G. Rubino, L. A. Rozema, A. Feix, M. Araújo, J. M. Zeuner, L. M. Procopio, Ā. Brukner and P. Walther. ‘Experimental Verification of an Indefinite Causal Order’ (2016). [arXiv:1608.01683 \[quant-ph\]](#).
- [17] G. Brassard, H. Buhrman, N. Linden, A. A. Méthot, A. Tapp and F. Unger. ‘Limit on Nonlocality in Any World in Which Communication Complexity Is Not Trivial’. *Phys. Rev. Lett.* **96** 250401 (2006). [eprint: quant-ph/0508042](#).

- [18] N. Linden, S. Popescu, A. J. Short and A. Winter. ‘Quantum Nonlocality and Beyond: Limits from Nonlocal Computation’. *Phys. Rev. Lett.* **99** 180502 (2007). [arXiv:quant-ph/0610097](#).
- [19] M. Pawłowski, T. Paterek, D. Kaszlikowski, V. Scarani, A. Winter and M. Żukowski. ‘Information causality as a physical principle’. *Nature* **461** 1101–1104 (2009). [arXiv:0905.2292 \[quant-ph\]](#).
- [20] M. Navascués and H. Wunderlich. ‘A glance beyond the quantum model’. *Proc. Royal Soc. A* **466** 881–890 (2009). [arXiv:0907.0372 \[quant-ph\]](#).
- [21] T. Fritz, A. B. Sainz, R. Augusiak, J. B. Brask, R. Chaves, A. Leverrier and A. Acín. ‘Local orthogonality as a multipartite principle for quantum correlations’. *Nat. Commun.* **4** 2263 (2013). [arXiv:1210.3018 \[quant-ph\]](#).
- [22] L. Hardy. ‘Quantum Theory From Five Reasonable Axioms’ (2001). [arXiv:quant-ph/0101012](#).
- [23] B. Dakić and Č. Brukner. ‘Deep Beauty: Understanding the Quantum World through Mathematical Innovation’. Cambridge University Press, 2011. Chap. Quantum Theory and Beyond: Is Entanglement Special?, pp. 365–392. [arXiv:0911.0695 \[quant-ph\]](#).
- [24] L. Masanes and M. P. Müller. ‘A derivation of quantum theory from physical requirements’. *New J. Phys.* **13** 063001 (2011). [arXiv:1004.1483 \[quant-ph\]](#).
- [25] G. Chiribella, G. M. D’Ariano and P. Perinotti. ‘Informational derivation of quantum theory’. *Phys. Rev. A* **84** 012311 (2011). [arXiv:1011.6451 \[quant-ph\]](#).
- [26] H. Barnum, M. P. Müller and C. Ududec. ‘Higher-order interference and single-system postulates characterizing quantum theory’. *New J. Phys.* **16** 123029 (2014). [arXiv:1403.4147 \[quant-ph\]](#).
- [27] P. A. Höhn. ‘Toolbox for reconstructing quantum theory from rules on information acquisition’ (2014). [arXiv:1412.8323 \[quant-ph\]](#).
- [28] P. A. Höhn and C. Wever. ‘Quantum theory from questions’ (2015). [arXiv:1511.01130 \[quant-ph\]](#).
- [29] A. Feix, M. Araújo and Č. Brukner. ‘Causally nonseparable processes admitting a causal model’. *New J. Phys.* **18** 083040 (2016). [arXiv:1604.03391 \[quant-ph\]](#).
- [30] G. Chiribella, G. M. D’Ariano and P. Perinotti. ‘Theoretical framework for quantum networks’. *Phys. Rev. A* **80** 022339 (2009). [arXiv:0904.4483 \[quant-ph\]](#).
- [31] G. C. Ghirardi, A. Rimini and T. Weber. ‘Unified dynamics for microscopic and macroscopic systems’. *Phys. Rev. D* **34** 470–491 (1986).
- [32] G. C. Ghirardi, P. Pearle and A. Rimini. ‘Markov processes in Hilbert space and continuous spontaneous localization of systems of identical particles’. *Phys. Rev. A* **42** 78–89 (1990).
- [33] A. Bassi, K. Lochan, S. Satin, T. P. Singh and H. Ulbricht. ‘Models of wave-function collapse, underlying theories, and experimental tests’. *Rev. Mod. Phys.* **85** 471–527 (2013). [arXiv:1204.4325 \[quant-ph\]](#).
- [34] S. W. Hawking. ‘Breakdown of predictability in gravitational collapse’. *Phys. Rev. D* **14** 2460–2473 (1976).
- [35] D. Harlow. ‘Jerusalem Lectures on Black Holes and Quantum Information’. *Rev. Mod. Phys.* **88** 015002 (2016). [arXiv:1409.1231 \[hep-th\]](#).
- [36] T. Jacobson. ‘Trans-Planckian Redshifts and the Substance of the Space-Time River’. *Progress of Theoretical Physics Supplement* **136** 1–17 (1999). [arXiv:hep-th/0001085](#).
- [37] M. Bojowald, D. Cartin and G. Khanna. ‘Lattice refining loop quantum cosmology, anisotropic models, and stability’. *Phys. Rev. D* **76** 064018 (2007). [arXiv:0704.1137 \[gr-qc\]](#).
- [38] S. Gielen and L. Sindoni. ‘Quantum Cosmology from Group Field Theory Condensates: a Review’. *SIGMA* **12** 082 (2016). [arXiv:1602.08104 \[gr-qc\]](#).
- [39] P. A. Höhn. ‘Quantization of systems with temporally varying discretization. I. Evolving Hilbert spaces’. *J. Math. Phys.* **55** 083508 (2014). [arXiv:1401.6062 \[gr-qc\]](#).
- [40] V. Mukhanov. *Physical Foundations of Cosmology*. Cambridge University Press, 2005. ISBN: 9780521563987.
- [41] K. Życzkowski and I. Bengtsson. *Geometry of Quantum States*. Cambridge University Press, 2006.
- [42] Č. Brukner. ‘Bounding quantum correlations with indefinite causal order’. *New J. Phys.* **17** 083034 (2015). [arXiv:1404.0721 \[quant-ph\]](#).
- [43] Ä. Baumeler and S. Wolf. Private communication (2015).
- [44] A. Royer. ‘Wigner function in Liouville space: A canonical formalism’. *Phys. Rev. A* **43** 44–56 (1991).
- [45] S. L. Braunstein, G. M. D’Ariano, G. J. Milburn and M. F. Sacchi. ‘Universal Teleportation with a Twist’. *Phys. Rev. Lett.* **84** 3486–3489 (2000). [arXiv:quant-ph/9908036](#).

A The Choi-Jamiołkowski isomorphism

We use two levels of the Choi-Jamiołkowski isomorphism. The first level is the “pure” CJ isomorphism, used to represent matrices as vectors. The CJ operator of a matrix $A : \mathcal{H}_I \rightarrow \mathcal{H}_O$ is its “double-ket” [44, 45]:

$$|A\rangle\rangle := \mathbb{1} \otimes A|\mathbb{1}\rangle\rangle = \sum_{i=0}^{d_{\mathcal{H}_I}-1} |i\rangle A|i\rangle \quad (30)$$

The second level is the “mixed” CJ isomorphism, used to represent linear operators that act on matrices as matrices themselves. The CJ operator of a map $\mathcal{M} : \mathcal{L}(\mathcal{H}_I) \rightarrow \mathcal{L}(\mathcal{H}_O)$ is

$$\mathfrak{C}(\mathcal{M}) := \mathcal{I} \otimes \mathcal{M}(|\mathbb{1}\rangle\rangle\langle\langle\mathbb{1}|) = \sum_{i,j=0}^{d_{\mathcal{H}_I}-1} |i\rangle\langle j| \otimes \mathcal{M}(|i\rangle\langle j|) \quad (31)$$

Note that this convention differs from the one in [1, 9] by a transposition. This is necessary to avoid representing completely positive maps with a non-positive matrix, as would happen with the convention from [1, 9] when, for example, we take in equation (1) the process as being identity from P to A_I and identity from A_O to F , and the map \mathcal{A} to be a SWAP.

B The link product

The motivation for defining the link product is to express the CJ operator of a composition of two linear maps \mathcal{M} and \mathcal{N} directly as a function of the CJ operators of the individual maps [30], as

$$\mathfrak{C}(\mathcal{N} \circ \mathcal{M}) = \mathfrak{C}(\mathcal{N}) * \mathfrak{C}(\mathcal{M}).$$

We do not need, however, to explicitly use the CJ isomorphism, as the link product can be more simply defined as an abstract operation on two matrices. Namely, the link product between two operators $M \in \bigotimes_{i \in I} A^i$ and $N \in \bigotimes_{i \in J} A^i$ is

$$N * M := \text{tr}_{I \cap J}[(\mathbb{1}^{J \setminus I} \otimes M^{T_{I \cap J}})(N \otimes \mathbb{1}^{I \setminus J})] \quad (32)$$

Note that when $I \cap J = \emptyset$ the link product reduces to $N * M = N \otimes M$ and, conversely, when $I \cap J = I \cup J$, it reduces to $N * M = \text{tr } M^T N$.

C Conditions for validity

In this appendix we derive the constraints on W such that the map

$$G_{A,B} = W * (A \otimes B)$$

is a valid CPTP map for all valid CPTP maps A and B . Recall that a linear map $\mathcal{A} : A_I \rightarrow A_O$ is completely positive and trace preserving if and only if its CJ operator $A = \mathfrak{C}(\mathcal{A})$ is positive and $\text{tr}_{A_O} A = \mathbb{1}^{A_I}$. Therefore we need that

$$\text{tr}_{A'_O B'_O F} W * (A \otimes B) = \mathbb{1}^{A'_I B'_I P} \quad \text{and} \quad W * (A \otimes B) \geq 0 \quad (33)$$

$$\forall A, B \quad \text{s.t.} \quad A \geq 0, \quad \text{tr}_{A_O A'_O} A = \mathbb{1}^{A_I A'_I} \quad (34)$$

$$B \geq 0, \quad \text{tr}_{B_O B'_O} B = \mathbb{1}^{B_I B'_I} \quad (35)$$

First we show that $W * (A \otimes B) \geq 0$ iff $W \geq 0$. To see that, note that when \mathcal{A} and \mathcal{B} are swaps we have $W * (A \otimes B) = W$, so $W \geq 0$ is a necessary condition for $G_{A,B} \geq 0$. To see that it is also sufficient, just note that the link product of two positive operators is always positive.

To get the constraints on W such that $\text{tr}_{A'_O B'_O F} W * (A \otimes B) = \mathbb{1}^{A'_I B'_I P}$, first note that the positivity of A, B , and W is not relevant for that, since the set of positive operators is a full dimensional subset of the space of hermitian operators. Therefore we only need the normalisation condition

$$\text{tr}_{A'_O B'_O F} W * (A \otimes B) = \mathbb{1}^{A'_I B'_I P} \quad (36)$$

$$\forall A, B \quad \text{s.t.} \quad \text{tr}_{A_O A'_O} A = \mathbb{1}^{A_I A'_I}, \quad \text{tr}_{B_O B'_O} B = \mathbb{1}^{B_I B'_I} \quad (37)$$

Note also that the dimension of A'_I and A'_O does not affect which set of operators $\text{tr}_{A'_I A'_O} A$ is⁷, so we can without loss of generality set $d_{A'_I} = d_{A'_O} = d_{B'_I} = d_{B'_O} = 1$, simplifying this condition to

$$\text{tr}_F W * (A \otimes B) = \mathbb{1}^P \quad (38)$$

$$\forall A, B \quad \text{s.t.} \quad \text{tr}_{A_O} A = \mathbb{1}^{A_I}, \quad \text{tr}_{B_O} B = \mathbb{1}^{B_I} \quad (39)$$

Note also that

$$\text{tr}_{A_O} A = \mathbb{1}^{A_I} \iff A = a - {}_{A_O} a + \frac{\mathbb{1}^{A_I A_O}}{d_{A_O}} \quad (40)$$

for some Hermitian operator a , where we are using the shorthand notation

$${}_X W = \frac{\mathbb{1}^X}{d_X} \otimes \text{tr}_X W. \quad (41)$$

Using this in Alice and Bob's sides allows us to reduce the normalization condition to

$$\forall a, b \quad (\text{tr}_F W) * \left[\left(a - {}_{A_O} a + \frac{\mathbb{1}^{A_I A_O}}{d_{A_O}} \right) \otimes \left(b - {}_{B_O} b + \frac{\mathbb{1}^{B_I B_O}}{d_{B_O}} \right) \right] = \mathbb{1}^P \quad (42)$$

For $a = b = 0$ this yields the condition

$$(\text{tr}_F W) * \left(\frac{\mathbb{1}^{A_I A_O}}{d_{A_O}} \otimes \frac{\mathbb{1}^{B_I B_O}}{d_{B_O}} \right) = \mathbb{1}^P \quad (43)$$

For $b = 0$ and $a = 0$, respectively, equation (43) together with (42) imply that

$$\forall a \quad (\text{tr}_F W) * \left[\left(a - {}_{A_O} a \right) \otimes \frac{\mathbb{1}^{B_I B_O}}{d_{B_O}} \right] = 0 \quad (44)$$

$$\forall b \quad (\text{tr}_F W) * \left[\frac{\mathbb{1}^{A_I A_O}}{d_{A_O}} \otimes \left(b - {}_{B_O} b \right) \right] = 0 \quad (45)$$

which then imply that

$$\forall a, b \quad (\text{tr}_F W) * [(a - {}_{A_O} a) \otimes (b - {}_{B_O} b)] = 0 \quad (46)$$

Using the fact that the maps ${}_{A_O}$ and ${}_{B_O}$ are self-adjoint under the Hilbert-Schmidt inner product, we see that conditions (44)-(46) are equivalent to

$$\text{tr}_{B_I B_O F} W = {}_{A_O} \text{tr}_{B_I B_O F} W \quad (47)$$

$$\text{tr}_{A_I A_O F} W = {}_{B_O} \text{tr}_{A_I A_O F} W \quad (48)$$

$$\text{tr}_F W = {}_{A_O} \text{tr}_F W + {}_{B_O} \text{tr}_F W - {}_{A_O B_O} \text{tr}_F W \quad (49)$$

These conditions together with (43) characterize completely the affine subspace to which valid process matrices belong. It is, however, convenient to rewrite them so that we characterize this affine subspace as a linear subspace together with a single constraint on the trace of W . To do that, note that conditions (47)-(49) define linear subspaces, so we only need to rewrite condition (43). It is easy to see that it is equivalent to:

$$\text{tr}_{A_I A_O B_I B_O F} W = \text{tr}_{P A_I A_O B_I B_O F} W \quad (50)$$

$$\text{tr} W = d_P d_{A_O} d_{B_O} \quad (51)$$

So the linear subspace of valid process matrices is the intersection of the linear subspaces defined in the equations (47)-(50). Since the projectors onto these subspaces commute, we can obtain the projector onto the subspace of valid process matrices L_V simply by composing them. Elementary manipulations give us then

$$\begin{aligned} L_V(W) = & W - {}_F W + {}_{A_O F} W + {}_{B_O F} W - {}_{A_O B_O F} W - {}_{A_I A_O F} W + {}_{A_I A_O B_O F} W \\ & - {}_{B_I B_O F} W + {}_{A_O B_I B_O F} W - {}_{A_I A_O B_I B_O F} W + {}_{P A_I A_O B_I B_O F} W \end{aligned} \quad (52)$$

To summarize the results of this section, W is a valid process matrix iff

$$W \geq 0 \quad (53)$$

$$\text{tr} W = d_P d_{A_O} d_{B_O} \quad (54)$$

$$W = L_V(W) \quad (55)$$

⁷It would if the CPTP maps A were restricted to be unitaries.

D The null-square lemma

The purpose of this section is to prove that, for any Hermitian matrix C , the greatest null square submatrix of C achievable via a change of basis has size $d_C \equiv \min(n_+, n_-) + n_0$, where n_+, n_-, n_0 denote, respectively, the dimensionality of the subspaces spanned by the eigenvectors of C with positive, negative and zero eigenvalues.

We will first establish that the above quantity is an upper bound for the greatest zero-square submatrix. Let then S be a subspace with the property that $\langle v|C|w\rangle = 0$, for $|v\rangle, |w\rangle \in S$, and denote by Π_0 the projector onto the null space of C .

Note that $\dim(S) \leq \dim(S_0) + \dim(S_{+-})$, where $S_0 \equiv \Pi_0 S$ and $S_{+-} \equiv (\mathbb{1} - \Pi_0)S$. Notice as well that $\langle v|C|w\rangle = 0$, for $|v\rangle, |w\rangle \in S_{+-}$. Clearly, the dimension of S_0 is upper bounded by n_0 , hence all we need to do is show that the dimension of S_{+-} is upper bounded by $\min(n_+, n_-)$.

Let $\{|u^{(i)}\rangle\}_i$ be a basis for S_{+-} , and let $\{\lambda_k\}_{k=1}^{n_+}$ ($\{-\mu_k\}_{k=1}^{n_-}$) be the set of positive (minus negative) eigenvalues of C , with eigenvectors $\{|\psi_k\rangle\}_{k=1}^{n_+}$ ($\{|\phi_k\rangle\}_{k=1}^{n_-}$). Then,

$$|u^{(i)}\rangle = \sum_{k=1}^{n_+} v_k^{(i)} |\psi_k\rangle + \sum_{k=1}^{n_-} w_k^{(i)} |\phi_k\rangle.$$

From the condition $\langle u^{(i)}|C|u^{(j)}\rangle = 0$ we have that

$$\sum_{k=1}^{n_+} (v_k^{(i)})^* v_k^{(j)} \lambda_k - \sum_{k=1}^{n_-} (w_k^{(i)})^* w_k^{(j)} \mu_k = 0,$$

or, in other words, $\langle \tilde{v}^{(i)}|\tilde{v}^{(j)}\rangle = \langle \tilde{w}^{(i)}|\tilde{w}^{(j)}\rangle$, with $\tilde{v}_k^{(i)} \equiv \sqrt{\lambda_k} v_k^{(i)}$, $\tilde{w}_k^{(i)} \equiv \sqrt{\mu_k} w_k^{(i)}$. This implies that there exists an isometry U such that $|\tilde{v}^{(i)}\rangle = U|\tilde{w}^{(i)}\rangle$. The vectors $|\tilde{v}^{(i)}\rangle \oplus |\tilde{w}^{(i)}\rangle$ (and thus $|u^{(i)}\rangle$) are generated by applying a linear transformation on $|\tilde{w}^{(i)}\rangle$ and so the space they span cannot be greater than n_- . An identical argument shows that the space spanned by $\{|u^{(i)}\rangle\}_{i=1}^m$ cannot be greater than n_+ either. We have just proven that d_C is an upper bound for the size of the null square.

To prove that it is also a lower bound, note that the $\min(n_+, n_-)$ linearly independent vectors $\{|u_i\rangle \equiv \lambda_k^{-1/2} |\psi_k\rangle + \mu_k^{-1/2} |\phi_k\rangle : i = 1, \dots, \min(n_+, n_-)\}$ satisfy $\langle u_i|C|u_j\rangle = 0$ for all i, j . Adding all n_0 vectors from the null space of C , we obtain a basis $\{|u_i\rangle\}_{i=1}^{d_C}$ with the property that $\langle u_i|A|u_j\rangle = 0$.